Heredity in Filum Disease: Arnold-Chiari I Syndrome, Idiopathic Syringomyelia and Idiopathic Scoliosis

Miguel B. Royo-Salvador

Institut Chiari & Siringomielia & Escoliosis de Barcelona

INTRODUCTION

Filum Disease encompasses a spectrum of conditions fundamentally driven by a mechanical conflict. This underlying conflict originates from the **asynchronous growth** between the **central nervous system** (neuroaxis) and the **vertebral column**, a developmental divergence observed across various species. In human embryogenesis, this mechanical tension can occur universally, resulting in clinical dysfunctions that are directly proportional to the intensity of the caudal traction exerted by the *filum terminale*.

Despite the clinical relevance of the core pathologies comprising Filum Disease, there remains a significant deficit of comprehensive genetic data. Current clinical evidence indicates a clear hereditary component, suggesting a potential autosomal recessive or autosomal dominant inheritance pattern. Consequently, future genomic research should be strategically directed toward identifying genetic alterations linked to posterior neuropore anomalies and caudal nervous system malformations. Crucially, investigations must target the specific embryological junction between the ectoderm and mesoderm, which is directly responsible for the formation of the *filum terminale.*

PRIMARY OBJECTIVES

1. Hereditary Factor Analysis: To conduct a comprehensive literature review identifying the hereditary factors implicated in the aetiopathogenesis of Filum Disease and its associated clinical spectrum.

2. Familial Aggregation Study: To analyse observational data from our institutional clinical cohort, specifically evaluating patients who share a confirmed diagnosis of Filum Disease with at least one other family member.

3. Genetic Susceptibility: To investigate and propose potential genetic loci that may predispose individuals to the onset and development of Filum Disease.

METHODOLOGY

1. LITERATURE REVIEW

Design: Comprehensive bibliographic review.

Databases: Scientific and medical search engines, primarily utilizing PubMed and Google Scholar, among other academic repositories.

2. CLINICAL OBSERVATIONAL STUDY & PATIENT SELECTION

Base Population: 950 patients treated at the Institut Chiari & Siringomielia & Escoliosis de Barcelona (ICSEB) between the years 1993 and 2015.

Inclusion Criteria: Extraction of patients diagnosed with Filum Disease presenting familial aggregation (minimum of two affected members within the same family unit).

Final Sample: 33 isolated family groups.

3. CLINICAL & RADIOLOGICAL ANALYSIS

Variables Assessed: Evaluation of phenotypic expression through clinical records and neuroimaging (MRI).

Comparative Pedigree Analysis: Detailed anatomo-clinical comparison within specific family units, with a special focus on monozygotic twins (Groups 1 and 19) to isolate biomechanical variables (caudal traction) from identical genetic baselines.


RESULTS

1. FAMILIAL AGGREGATION IN FILUM DISEASE

Clinical Cohort: Out of a primary population of 950 patients treated for Filum Disease (1993–2015), 33 distinct family groups were isolated.

Hereditary Pattern: Each identified group comprises at least two affected members, empirically substantiating a robust hereditary substrate in the clinical manifestation of this pathological spectrum.


STUDY POPULATION



2. PHENOTYPIC DISCORDANCE IN MONOZYGOTIC TWINS

Anatomo-clinical Correlation (Family Group 1): Despite sharing identical genomes, the twins exhibit significant phenotypic discordance. Twin A presents a strictly caudal topography of the conus medullaris (L1-L2), which directly correlates with amplified neuroaxis traction and a significantly larger **syringomyelic cavity** volume compared to Twin B, whose conus medullaris is located more cranially.

Biomechanical Accommodation (Family Group 19): Identical genomes exhibit variable phenotypic expression dictated by mechanical traction. The manifestation of **idiopathic scoliosis** acts as a compensatory mechanism, distributing caudal traction and effectively preventing the centromedullary ischemia (syringomyelia) observed in the co-twin.



3. SYNDROMIC ASSOCIATIONS AND METHODOLOGICAL BIAS


Genetic Concurrence: Filum Disease frequently co-occurs with established genetic entities, including Klippel-Feil syndrome, Ehlers-Danlos syndrome, and von Recklinghausen neurofibromatosis.

Paradigm Shift: Contemporary genetic research is constrained by a methodological bias, erroneously focusing on secondary anatomical consequences such as posterior cranial fossa malformations or altered cerebrospinal fluid dynamics. Valid aetiological research must reorient towards tissue alterations at the specific ectoderm-mesoderm junction comprising the *filum terminale*.


METHODOLOGICAL BIAS



CAUDAL TRACTION



DISCUSSION



A Complex Inheritance Model

The pathologies comprising Filum Disease do not follow a simple Mendelian transmission. The clinical manifestation is determined by an oligogenic or **polygenic** architecture interacting with **environmental** factors, which modulate both the **phenotype** and **disease progression**.

The Monozygotic Paradox

While identical twins exhibit higher diagnostic concordance than dizygotic twins, they frequently present marked discordance in clinical severity, age of onset, and the presence of syringomyelia (SM). This phenotypic variability within identical genomes confirms that genetic predisposition alone is insufficient to explain the full anatomical spectrum.


Mesodermal Origin & Syndromic Co-inheritance

The frequent co-segregation of Chiari Malformation Type I (CMI) with bone and connective tissue disorders (such as Klippel-Feil, Goldenhar, and Ehlers-Danlos syndromes) strongly supports a mesodermal origin, with potential pleiotropic effects linked to chromosomes 9 and 15.

The Biomechanical Core

The fundamental cause of Filum Disease is the mechanical conflict initiated during the ninth embryonic week. Therefore, what is genetically inherited is not the pathology itself, but rather the **intensity of the asynchronous growth** between the neuroaxis and the vertebral column, combined with the histological resistance of the *filum terminale*.

THE BIOMECHANICAL CORE



CONCLUSIONS

GENETIC SUBSTRATE: Filum Disease exhibits a clear hereditary basis, taxonomically hypothesised as a congenital entity with a complex, potentially polygenic, and primarily autosomal recessive architecture.

PHENOTYPIC EXPRESSION: The clinical manifestation shows a marked female predominance and results from a complex interplay between multigenic alterations and environmental, chemical, or physical factors.

METHODOLOGICAL LIMITATIONS: Conventional genetic studies are limited in validity, having historically focused on derived anatomical consequences (e.g., craniocervical malformations, altered CSF dynamics) instead of isolating the primary aetiology.

FOUNDATIONAL EVIDENCE: Intrafamilial clinical analyses and the phenotypic discordance documented in monozygotic twins provide robust evidence for a foundational genetic basis underlying this pathology.

CLINICAL IMPLICATIONS AND RECOMMENDATIONS

Epidemiological Impact & Familial Risk: Clinical observations suggest a complex genetic architecture, potentially following an autosomal recessive or oligogenic inheritance pattern. Despite this hypothesised mechanism, Filum Disease exhibits a significantly high estimated global incidence. Crucially, familial aggregation indicates that descendants of affected individuals may face an increased probability (up to 65%) of developing the condition, underscoring the need for proactive familial communication.

Diagnostic Screening: Early intrafamilial screening is strongly recommended to monitor congenital progression. Initial preventative assessments should include a scoliogram to identify subtle idiopathic deviations, followed by a comprehensive cranio-vertebral MRI if suggestive symptomatology arises.

Differential Diagnosis: Due to its plurisyndromic clinical presentation, Filum Disease is frequently misdiagnosed as fibromyalgia, multiple sclerosis, or functional psychiatric disorders. Recognising the underlying biomechanical aetiology is essential to prevent misdirected therapeutic approaches.

REFERENCES

- Boyles AL,. Enterline DS., Hammock PH., Siegel DG., Slifer SH., Mehltretter L., Gilbert JR., Hu-Lince D., Stephan D., Batzdorf U., Benzel E., Ellenbogen R., Green BA., Kula R., Menezes A., Mueller D., Oro' JJ., Iskandar BJ., George TM., Milhorat TH., Speer MC. 2006. Phenotypic definition of Chiari type I malformation coupled with high-density SNP genome screen shows significant evidence for linkage to regions on chromosomes 9 and 15. *Am J Med Genet A*, 140(24):2776-2785.
- Turgut M (2001) Chiari type I malformation in two monozygotic twins. Brit J Neurosurg 15:279–280
- Waller T., Nowak R., Tkacz M., Zapart D., Mazurek U. 2013. Familiar or sporadic Idiopathic Scoliosis – classification based on artificial neural network and GAPDH and ACTB transcription profile. *BioMedical Engineering OnLine* 12:1-14.
- Milhorat TH, MD, Chou MW, MD, Trinidad EM, MD, Kula RW, MD, Mandell M, MD, Wolpert Chantelle, M.B.A. PA-C, Speer MC, PD (1999) Chiari I Malformation Redefined: Clinical and Radiographic Findings for 364 Symptomatic Patients. Neurosurgery 44:1005–1017 PMID: 10232534
- Milhorat TH, Bolognese PA, Nishikawa M, McDonnell NB, Francomano CA (2007) Syndrome of occipitoatlantoaxial hypermobility, cranial settling, and chiari malformation type I in patients with hereditary disorders of connective tissue. J Neurosurg Spine 7(6):601–609

Dr. Miguel B. Royo-Salvador
mroyo@institutchiaribcn.com